

Existence of $\sigma(600)/\kappa(900)$ -Particle and New Chiral Scalar Nonet “Chiralons”

Muneyuki Ishida* and Shin Ishida†

* *Department of Physics, University of Tokyo, Tokyo 113 and † Atomic Energy Research Institute, College of Science and Technology, Nihon University, Tokyo 101, JAPAN*

Abstract. The $\sigma(600)$ and $\kappa(900)$, observed in the phase shift analyses [1,2], satisfy rather well the mass and width relation predicted by the $SU(3)L\sigma M$ and the $SU(3)L\sigma M$ with the vector and axial-vector meson nonets, and deserve to be the members of scalar σ -nonet, together with the observed resonances $a_0(980)$ and $f_0(980)$, as a chiral partner of pseudoscalar π -nonet. In the phase shift analyses an introduction of repulsive background phase δ_{BG} is essential, whose origin has a close connection to the $\lambda\phi^4$ interaction in $L\sigma M$. It is argued that the members of this σ -nonet, “Chiralons,” have different properties and should be discriminated from the conventional 3P_0 - $q\bar{q}$ -scalar nonet.

(*Properties of σ/κ and Linear σ Model*) The $\sigma(600)$ and $\kappa(900)$ are observed by the reanalyses [1,2] of $\pi\pi$ and $K\pi$ phase shifts, respectively. Here their properties in relation to the $SU(3)$ chiral symmetry is examined. In the low energy region, where the structure of composite meson can be neglected, the linear σ model ($L\sigma M$) may be valid. In terms of matrix notation $B \equiv \sigma + i\phi$ ($\sigma \equiv \lambda^i \sigma^i / \sqrt{2}$ and $\phi \equiv \lambda^i \phi^i / \sqrt{2}$ denoting scalar and pseudoscalar meson nonet, respectively), the Lagrangian of $SU(3)L\sigma M$ [5,6] is

$$\begin{aligned} \mathcal{L} = & \frac{1}{2} \langle \partial_\mu B \partial^\mu B^\dagger \rangle - \frac{\mu^2}{2} \langle BB^\dagger \rangle - \frac{\lambda_1}{4} \langle BB^\dagger \rangle^2 - \frac{\lambda_2}{2} \langle (BB^\dagger)^2 \rangle \\ & + \kappa_d (\det B + \det B^\dagger) + \langle f\sigma \rangle, \end{aligned} \quad (1)$$

where $\langle \rangle$ represents the trace, and the form of $f (= \text{diag}\{f_n, f_n, f_s\})$ guarantees the PCAC. The six model parameters contained in Eq.(1) are determined by the masses of π, η, η', σ and κ , and the decay constant f_π , and then we can predict the properties of scalar mesons, given in Table 1: ¹ The predicted widths of σ and κ are consistent to the experimental values. Thus $\sigma(600)$ and $\kappa(900)$ have plausible

¹⁾ The predicted properties of scalar mesons are very sensitive to the value of f_K/f_π [6]. I take the region of this ratio $1.329 < f_K/f_\pi < 1.432$, where the m_κ^{theor} reproduces the experimental value within its uncertainty, being somewhat larger than the experimental one, $f_K/f_\pi = 113/93 = 1.22$.

TABLE 1. The masses and widths of scalar meson nonet(“chiralons”) predicted by SU(3)L σ M, compared with experiments (the unit is in MeV). The values with underlines are used as inputs. The $\Gamma_{\sigma'}^{\text{theor}}$ is the $\pi\pi$ -width.

	m^{theor}	m^{exp}	Γ^{theor}	Γ^{exp}		m^{theor}	m^{exp}	Γ^{theor}	Γ^{exp}
σ	<u>535~650</u>	<u>535~650</u>	400~800	385 $\pm$ 70	δ	900~930	984 $\pm$ 1	110~170	50~100
κ	<u>905$^{+65}_{-30}$</u>	<u>905$^{+65}_{-30}$</u>	300~600	545 $^{+235}_{-110}$	σ'	1030~1200	980 $\pm$ 10	0~300	40~100




FIGURE 1. $\pi\pi$ and $K\pi$ scattering diagram in L σ M

properties as the members of the scalar nonet in L σ M. The predicted properties of the other members, $\delta(I=1)$ and $\sigma'(s\bar{s})$, are close to those of $a_0(980)$ and $f_0(980)$. Thus, it may be plausible to regard **the $\sigma(600)$, $\kappa(900)$, $a_0(980)$, and $f_0(980)$ as being the members of the scalar nonet forming with the members of π -nonet the linear representation of the SU(3) chiral symmetry.**

(*repulsive core and $\lambda\phi^4$ term*) In our phase shift analyses the repulsive background phase shift of hard core type δ_{BG} introduced phenomenologically plays an essential role. The origin of this δ_{BG} seems to have a close connection to the $\lambda\phi^4$ -interaction in L σ M [4]: It represents a contact zero-range interaction and is strongly repulsive, and has plausible properties as the origin of repulsive core. In the following we shall make a tentative theoretical estimate of the core radii in the framework of L σ M: The diagrams for the $\pi\pi$ and $K\pi$ scattering amplitudes in the tree level, are given in Fig. 1. The scattering $\mathcal{T}$ matrix consists of a resonance part $\mathcal{T}_R$ and of a background part $\mathcal{T}_{BG}$. The $\mathcal{T}_{BG}$ corresponds to the contact $\lambda\phi^4$ interaction and the exchange of relevant resonances. This $\mathcal{T}_{BG}$ has a weak s -dependence in comparison with that of $\mathcal{T}_R$. We regard them as $\mathcal{K}$ -matrix ($\mathcal{K}_R \equiv \mathcal{T}_R$, $\mathcal{K}_{BG} \equiv \mathcal{T}_{BG}$), and unitarize them separately by $S = \prod_A S_A$ [2]; $S_A = (1 + i\rho_1\mathcal{K}_A)/(1 - i\rho_1\mathcal{K}_A)$ ($A = R, BG$). We estimate the theoretical core radius at the resonance energy by the formula, $r_c^{\text{theor}} \equiv \frac{1}{p_1} \text{atan}(-\rho_1\mathcal{T}_{BG})|_{\sqrt{s}=m_R}$, and the results are given in Table 2. They seem almost consistent to the experimental values obtained in the phase shift analysis [2].

(*Why σ and κ have been overlooked*) According to the above analysis the $\sigma(\sigma')$ consists almost of $n\bar{n}$ -($s\bar{s}$)-component, and I take the SU(2)L σ M for simplicity. The

TABLE 2. Repulsive core radii $r_c(\text{fm})$ in $\pi\pi$ and $K\pi$ scattering

	$(I=0)_{\pi\pi}$	$(I=1/2)_{K\pi}$	$(I=2)_{\pi\pi}$	$(I=3/2)_{K\pi}$
experiment	0.60 ± 0.07	0.70 ± 0.08	0.17	0.16
theory(L σ M)	$0.6 \sim 0.8$	$0.3 \sim 0.4$	$0.13 \sim 0.17$	$0.11 \sim 0.15$

$\pi\pi$ -scattering $A(s, t, u)$ -amplitude is given by $A(s, t, u) = (-2g_{\sigma\pi\pi})^2/(m_\sigma^2 - s) - 2\lambda$. Because of the relation, $g_{\sigma\pi\pi} = f_\pi\lambda = (m_\sigma^2 - m_\pi^2)/(2f_\pi)$, the amplitude due to virtual σ production(1st term) is exactly cancelled by that due to repulsive $\lambda\phi^4$ interaction(2nd term) in $O(p^0)$ level, and the $A(s, t, u)$ is rewritten into the following form:

$$A(s, t, u) = \frac{1}{f_\pi^2} \frac{(m_\sigma^2 - m_\pi^2)^2}{m_\sigma^2 - s} - \frac{m_\sigma^2 - m_\pi^2}{f_\pi^2} = \frac{s - m_\pi^2}{f_\pi^2} + \frac{1}{f_\pi^2} \frac{(m_\pi^2 - s)^2}{m_\sigma^2 - s}, \quad (2)$$

where in the last side the $O(p^2)$ Tomozawa-Weinberg amplitude and the $O(p^4)$ (and higher order) correction term are left. As a result the derivative coupling property of π -meson as a Nambu-Goldstone boson is preserved. In this sense the $\lambda\phi^4$ -interaction can be called a “compensating” interaction for σ -effect.

The essential point in our phase shift analyses is a strong cancellation between the positive δ_σ (δ_κ) and the negative δ_{BG} , as shown in Table 1 in the contribution to this conference [1], which corresponds to the Eq.(2) in L σ M.

In the conventional Breit-Wigner formula a non-derivative $\sigma\pi\pi$ -coupling, accordingly a non-derivative effective $\pi\pi$ -interaction, is supposed without consideration on the “compensating” repulsive interaction. This seems [4] to be a reason why σ (and κ) has been overlooked in most of the $\pi\pi$ [8] (and $K\pi$) phase shift analyses.

(Related problems) An investigation of the properties of the scalar σ -nonet with a wide scope is done by using the “super” Lagrangian [5], where the vector(ρ -) and the axial-vector(a_1 -) meson nonets are included consistently to the vector meson dominance. The eight new parameters in addition to the six parameters in Eq.(1) are appeared in this scheme. In Table 3, the theoretical values of 21 quantities(including f_π and f_K) are given in comparison with the experimental values. The obtained values of scalar meson properties are quite close to those in the original L σ M, Eq.(1). Thus now it becomes more plausible than before that the $\sigma(600)$, $\kappa(900)$, $\delta(980)(=a_0(980))$ and $\sigma'(980)(=f_0(980))$ form the chiral scalar-nonnet, which was first suggested in ref. [7].

It should be noted that the σ is now directly observed in the production processes as shown in the contribution to this conference [3]. Thus, the conventional argument [9] against the validity of L σ M and the existence of σ -meson in the framework of chiral perturbation theory(ChPT), which is based only on the indirect experiments concerning a much lower energy region than m_σ , is not appropriate. The parameters in L σ M are determined through the physics in resonance energy region, and the related low energy quantities are predicted with no free fitting parameters, quite

TABLE 3. Properties of scalar, pseudoscalar and vector mesons by the L σ M with ρ - and a_1 -nonets. (a) Masses and widths(MeV) of scalar mesons. (b) Masses(MeV) of pseudoscalar and vector mesons and widths(MeV) of vector mesons. The $f_\pi(f_K)$ is obtained with 93.2(114.7)MeV, consistent to the experimental value .

(a)	m_σ	m_κ	m_δ	$m_{\sigma'}$	Γ_σ	Γ_κ	Γ_δ	$\Gamma_{\sigma'}$
theor	600	989	955	1063	454	505	174	131
exp	535~675	905^{+65}_{-30}	984 ± 1	980 ± 10	385 ± 70	545^{+235}_{-110}	50~100	40~100

(b)	m_π	m_K	m_η	$m_{\eta'}$	m_ρ	m_{K^*}	m_ω	m_ϕ	Γ_ρ	Γ_{K^*}	Γ_ϕ
theor	139	502	559	1020	783	887	778	1021	150	57	3
exp	138	495	547	958	769	894	782	1019	151	50	4

in contrast with the situation in ChPT. For example, the $\pi\pi$ -scattering lengths a_J^I obtained by the L σ M with ρ - and a_1 -nonets are $(a_0^0, a_0^2; a_1^1; a_2^0, a_2^2; a_3^1) = (0.16, -0.010; 0.022; 13 \times 10^{-4}, 2.3 \times 10^{-4}, 4 \times 10^{-5})$, which are not so much different from the present experimental values . In this connection we should like to refer to the work [10] where the width and form factor of K_{l4} -decay, as well as the $\pi\pi$ -phase shift δ_0^0 in low energy region, are also well described by L σ M similarly as in ChPT.²

Finally it is also to be noted that the members of the σ -nonet (the a_1 -nonet), “Chiralons,” have different properties and should be discriminated from the conventional 3P_0 -(3P_1 -) $q\bar{q}$ -scalar nonet. (See, the contribution [1].)

REFERENCES

1. S. Ishida et al., this conference; *Prog. Theor. Phys.***95**(1996)745;**98**(1997)1005;621.
2. T. Ishida et al., this conference.
3. K. Takamatsu et al., this conference. T. Ishida, Doctor thesis, Univ. of Tokyo 1996.
4. M.Y.Ishida, et al., this conference. M.Y.Ishida, *Prog. Theor. Phys.***96**(1996)853.
5. G. Gasiorowicz and D. A. Geffen, *Rev. Mod. Phys.* **3**(1969)531.
6. L.H.Chan and R.W.Haymaker, *Phys.Rev.***D7**(1973)402;**10**(1974)4143;4170.
7. M. D. Scadron, *Phys. Rev.* **D26**, 239 (1982).
8. K. L. Au, D. Morgan and M. R. Pennington, *Phys. Rev.* **D41**, 1633 (1987).
9. J.Gasser and H.Leutwyler, *Ann.Phys.***158**(1984)142; *Nucl.Phys.***B250**(1985)465;517.
10. E. P. Shabalin, *Yadern. Fiz.* **49**, 588 (1989) (*Sov. J. Nucl. Phys.* **49**, 365 (1989)).
11. U. G. Meissner, *Comm. Nucl. Part. Phys.***20**(1991)119.

²⁾ It is discussed [11] that the effect of σ -meson can be taken into account through the $O(p^4)$ and $O(p^6)$ Lagrangian in ChPT. Such a non-linear approach has no predictive power about the properties of the scalar σ -meson nonet as a chiral partner of the π -nonet, such as shown in Tables 1 and 3.